PART 4 OF 6

ARRT Trap-Buster: Radiation Protection & Radiobiology

High-Yield Dose Limits, Fluoroscopy Standards, Shielding Rules, Cell Sensitivity & Acute/Fetal Syndromes

1. NCRP Report #116 Annual & Cumulative Dose Limits

Registry Rule: NCRP limits apply to *occupational and public exposure*—they **never** include natural background radiation (~3.1 mSv/yr) or a patient's own medical/dental X-ray exams.

Exposed Group / Organ	SI Limit	Conventional Limit	Common ARRT Exam Trap
Occupational — Annual Effective Dose (EfD)	**50 mSv / yr**	5 rem / yr	Often confused with 500 mSv (skin/hands). Whole-body occupational cap is **50 mSv**.
Occupational — Cumulative Lifetime Limit	**10 mSv × Age**	1 rem × Age	Uses **10 mSv**, NOT 50 mSv (e.g., 28-yr-old tech = 280 mSv cumulative limit).
Lens of the Eye (Annual EqD)	**150 mSv / yr**	15 rem / yr	Separate from whole body (50) and extremities (500). Cataracts are deterministic.
Skin, Hands & Feet (Annual EqD)	**500 mSv / yr**	50 rem / yr	Highest single-organ occupational limit; applies to localized extremity exposure.
Embryo / Fetus — Total Gestation	**5 mSv**	0.5 rem	Applies **only after pregnancy is declared** in writing to the employer.
Embryo / Fetus — Monthly Limit	**0.5 mSv / mo**	0.05 rem / mo	Cannot exceed **0.5 mSv in any single month**, even if total gestation is under 5 mSv.
General Public — Frequent / Continuous	**1 mSv / yr**	0.1 rem / yr	Applies to waiting rooms, hallways, and uncontrolled areas outside X-ray rooms.
General Public — Infrequent Exposure	**5 mSv / yr**	0.5 rem / yr	Applies to occasional visitors or non-occupational individuals holding a patient.
Minors / Students Under Age 18	**1 mSv / yr**	0.1 rem / yr	Trainees under 18 are capped at the general public frequent limit (1 mSv/yr).

TRAP ALERT: Dosimeter Placement & Declared Pregnancy Rules

- **Single Badge Location:** Worn at the **collar level** outside the lead apron (anteriorly) to estimate head/neck and lens of eye dose.
- **Two-Badge System (Fluoro / Interventional):** Badge #1 at the **collar outside** the apron; Badge #2 at the **waist underneath** the apron.
- **Pregnant Radiographer:** Wears a dedicated fetal monitor at the **waist level under the lead apron** (never outside the apron).
- **Minimum Reporting Level:** OSL dosimeters detect down to **0.01 mSv (1 mrem)**; TLD down to **0.05 mSv (5 mrem)**; film badges down to **0.10 mSv (10 mrem)**. Readings below threshold report as "M" (Minimal).

2. Fluoroscopy, Mobile & Structural Shielding Standards

Equipment / Barrier Standard	Mandatory Limit / Spec	Registry Trap & Clinical Reasoning
Standard Fluoro Tabletop Rate	**≤ 100 mGy/min** (10 R/min)	Measured at the tabletop (or 30 cm in front of image intensifier on C-arms).
High-Level Control (HLC) Fluoro	**≤ 200 mGy/min** (20 R/min)	Requires continuous manual activation and a continuous audible signal while active.
Fluoro Cumulative Timer	**5 minutes** (300 seconds)	Sounds an audible alarm at 5 minutes of beam-on time; **does NOT shut off** the beam.
Minimum Source-to-Skin Distance	**38 cm (15 in)** stationary **30 cm (12 in)** mobile C-arm	Prevents excessive skin entrance dose from inverse square law proximity. Spacer locks enforce this.
Bucky Slot Cover & Fluoro Curtain	**0.25 mm Pb** equivalent	Blocks scatter emerging from the Bucky slot opening and the side of the tower at gonadal/waist level.
Protective Lead Aprons & Gloves	**0.5 mm Pb** (standard) **0.25 mm Pb** (minimum)	0.5 mm Pb attenuates ~90% of scatter at 75 kVp; **0.25 mm Pb** is minimum for thyroid/gloves.
Primary Protective Barrier	**1/16 inch (1.5 mm) Pb**	Stops the **useful primary beam**; extends **7 feet (2.1 m)** high on walls hit directly.
Secondary Protective Barrier	**1/32 inch (0.8 mm) Pb**	Stops **scatter and leakage radiation only**; includes control booth barrier and upper walls (>7 ft).
Control Booth Viewing Window	**1.5 mm Pb** equivalent	Constructed of lead-impregnated glass/acrylic; exposure switch cord must be too short to reach outside booth.
Mobile Exposure Cord & Stand	**≥ 6 feet (2 meters)** cord	Tech stands at a **90° angle** to the patient and at least 6 feet away (lowest scatter zone).

CARDINAL PRINCIPLES & BARRIER DESIGN FACTORS

- **Three Cardinal Principles:** Minimize **Time**, maximize **Distance** (most effective!), and use **Shielding**.
- **Four Structural Shielding Factors: Workload (W)** in mA-min/week, **Use Factor (U)** (fraction of time beam hits a primary wall; U = 1 for secondary barriers), **Occupancy Factor (T)** (1 for full-time offices, 1/4 for corridors, 1/16 for restrooms/stairways), and **Distance (d)**.

PART 4 OF 6

Radiobiology: Cell Sensitivity, ARS & Fetal Effects

Bergonié-Tribondeau Law, Target Theory, Dose-Response Curves & Deterministic vs. Stochastic Effects

3. Cellular Radiosensitivity & Radiobiology Laws

LAW OF BERGONIÉ AND TRIBONDEAU (1906)

Radiosensitivity of cells is **directly proportional** to their reproductive activity (mitotic rate) and **inversely proportional** to their degree of differentiation (maturity). Young, rapidly dividing, undifferentiated stem cells are the easiest to damage.

Sensitivity Level	Specific Cell Types	Registry Exam Traps
Highest Radiosensitivity	**Lymphocytes**, Spermatogonia, Erythroblasts, Intestinal Crypt Cells	**Lymphocytes** (white blood cells) are the **most radiosensitive** cells in the human body—even though they do not divide!
Intermediate Sensitivity	Endothelial cells, Osteoblasts, Spermatids, Fibroblasts	More sensitive than mature bone/muscle, less sensitive than blood stem cells.
Lowest Radiosensitivity (Radioresistant)	**Nerve cells (neurons)**, **Muscle cells (myocytes)**, Mature Red Blood Cells (RBCs)	Highly differentiated and non-dividing; mature RBCs lack a nucleus and are very radioresistant.

- **Direct vs. Indirect Action: Direct action** occurs when radiation hits DNA directly (dominant with high-LET radiation like alpha particles). **Indirect action** occurs when radiation ionizes water molecules (**radiolysis of water**), creating toxic **free radicals (H• and OH•)** that damage DNA—accounts for **~80% to 95%** of low-LET X-ray biological damage.
- **Oxygen Enhancement Ratio (OER):** Oxygenated tissue is **2.5 to 3 times more sensitive** to low-LET X-rays than hypoxic (oxygen-starved) tissue because oxygen locks in free radical damage (peroxide formation). For high-LET radiation, OER = 1.
- **LET vs. RBE: Linear Energy Transfer (LET)** = energy deposited per unit track length (keV/μm). X-rays are **low-LET (~3 keV/μm)**. As LET increases, **Relative Biological Effectiveness (RBE)** increases up to ~100 keV/μm, then drops due to "overkill" wasted energy.
- **Cell Cycle Sensitivity:** Most radiosensitive during **M phase (Mitosis)** and **G2 phase**; most radioresistant during **late S phase (DNA synthesis)**.

4. Stochastic vs. Deterministic Effects & Dose-Response Curves

Feature	Stochastic (Probabilistic) Effects	Deterministic (Tissue Reaction) Effects
Threshold Dose	**NO threshold** (any dose carries some risk)	**YES — clear threshold** must be exceeded
Dose Relationship	**Linear-Nonthreshold (LNT)** model	**Linear-Threshold** or **Nonlinear-Threshold (Sigmoid)**
Effect of Higher Dose	Increases **PROBABILITY** of occurrence, NOT severity	Increases **SEVERITY** of the injury
Classic Examples	**Radiation-induced cancer** (leukemia, thyroid, breast) and **genetic/hereditary mutations**	**Skin erythema** (2 Gy threshold), epilation (hair loss, 3 Gy), **cataracts** (2 Gy single / 5 Gy protracted), sterility

5. Acute Radiation Syndrome (ARS) & Fetal Gestation Windows

Syndrome / Stage	Whole-Body Threshold	Clinical Mechanism & Survival Timeline
Prodromal Stage (ARS)	**≥ 1 Gy** (100 rad)	Immediate nausea, vomiting, diarrhea (NVD) lasting hours to 2 days, followed by a symptom-free **Latent Period**.
Hematologic Syndrome	**1 – 10 Gy** (100–1,000 rad)	Destroys bone marrow stem cells → infection, hemorrhage, anemia. Death in **30–60 days**; **$LD_{50/60}$** for humans is **3.5 – 4.5 Gy** without medical care.
Gastrointestinal (GI) Syndrome	**6 – 10 Gy** (600–1,000 rad)	Destroys intestinal epithelial lining → severe bloody diarrhea, dehydration, electrolyte collapse. Death in **3 – 10 days**.
Central Nervous System (CNS)	**> 50 Gy** (> 5,000 rad)	Vasculitis, cerebral edema, intracranial pressure, seizures, coma. Death within **hours to 3 days** (no latent period).

TRAP ALERT: In-Utero Fetal Irradiation Windows (Must Know All Three!)

- **1. Preimplantation (0 – 2 Weeks post-conception):** Follows the **"All-or-Nothing" rule**. High doses (~0.1–0.2 Gy) cause embryonic resorption/death; surviving embryos develop normally without congenital malformations.
- **2. Major Organogenesis (2 – 8 Weeks post-conception): MOST SENSITIVE stage for congenital abnormalities** (skeletal defects, microcephaly, cleft palate, organ malformation) because organs are actively differentiating.
- **3. Fetal Growth Stage (8 Weeks – Birth):** Lower risk of gross structural defects, but highest risk for **growth retardation, intellectual disability (mental retardation)**, and **childhood leukemia/cancer** (10-Day Rule / 28-Day Rule context).